

Intra-ovarian injection of bone marrow-derived c-Kit⁺ cells for ovarian rejuvenation in menopausal rats

Sepideh Sheshpari^{1#}, Mahnaz Shahnazi¹, Shahin Ahmadian^{2#}, Mohammad Nouri^{3,4}, Mehran Mesgari Abbasi⁵, Rahim Beheshti⁶, Reza Rahbarghazi^{3,7}, Ali Honaramooz⁸, Mahdi Mahdipour^{3,4*}

¹ Department of Midwifery, Faculty of Nursing and Midwifery, Tabriz University of Medical Sciences, Tabriz, Iran

² Department of Biology, Faculty of Science, Azerbaijan Shahid Madani University, Tabriz, Iran

³ Stem Cell Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

⁴ Department of Reproductive Biology, Faculty of Advanced Medical Sciences, Tabriz University of Medical Sciences, Tabriz, Iran

⁵ Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

⁶ Department of Veterinary Science, Islamic Azad University, Shabestar Branch, Shabestar, Iran

⁷ Department of Applied Cell Sciences, Faculty of Advanced Medical Sciences, Tabriz University of Medical Sciences, Tabriz, Iran

⁸ Department of Veterinary Biomedical Sciences, Western College of Veterinary Medicine, University of Saskatchewan, Saskatoon, SK, S7N 5B4, Canada




Fig. S1. Body weight of rat's pre-transplantation (every 5 days) and post-transplantation (weekly) in different groups.